



Monoallelic STAT1 mutations and disease patterns

ABSTRACT

Monoallelic mutations in STAT1 are associated with a myriad of clinical phenotypes. Some clinical patterns appear to be preferentially associated with mutations in various STAT1 domains. Included are trends of phenotype-genotype correlations in patients with mutations in STAT1.

In the past several years it became apparent that mutations in *STAT1* are the most common cause of chronic mucocutaneous candidiasis (CMCC) in North America.

The clinical features of thrush and skin and nail lesions are indistinguishable between CMCC caused by an autosomal recessive mutation in the *AIRE* gene, which was described 2 decades ago (Ahonen et al. 1990; Finnish-German APECED Consortium 1997), or the recently identified monoallelic mutations in *STAT1* (van de Veerdonk et al. 2011) (Figure 1). Autoimmune manifestations including endocrinopathies are also shared between these genetic entities. However, it appears that although hypoparathyroidism and hypoadrenodism are more common in *AIRE* deficiency, hypothyroidism and diabetes mellitus are more common with *STAT1* mutations. Moreover, recurrent respiratory infections and antibody deficiency have a confirmed association with STAT1 abnormalities, but only rarely in *AIRE* deficiency (Bentur et al. 1991).

Interestingly, clinical entities formerly thought to be distinct, such as autosomal dominant CMCC, are also associated with mutations in *STAT1*. The clinical spectrum in these cases is wide (see article by Nahum and Dalal on page 97) ranging from vascular disease in the central nervous system to very mild symptoms of isolated oral thrush. Upon review of all published cases (Liu et al. 2011; Maródi et al. 2012; Takezaki et al. 2012; Sampaio et al. 2013; Uzel et al. 2013; Sharfe et al. 2014), it appears that mutations associated with autosomal

dominant CMCC are mostly localized to the coiled-coil domain of *STAT1*. The mean and median ages of both of these patients at the time of reporting appears to be around 30 years of age (Figure 2), but more striking are the mean and median ages of death in this group of patients as young as 35 years and 28 years, respectively (Figure 3). The cause of death was squamous cell carcinoma in the older patients and stroke in younger individuals.

Recently, whole exome sequencing discovered a surprising phenotype of monoallelic *STAT1* deficiency (Sharfe et al. 2014). Invariably, this group of patients had no family history of immunodeficiency. They present early in life with repeated viral and bacterial infections, and they suffer chronic colitis as well as other autoimmune manifestations. Evaluation of their immune system reveals a progressive decline in immunity, culminating in a combined immunodeficiency including profound lymphopenia and antibody deficiency, practically developing combined immunodeficiency (CID). Most of these patients have mutations in the DNA binding domain (DBD) of *STAT1*.

These patients with sporadic CID invariably die of overwhelming viral infection at a mean and median age of 10 years. This analysis, albeit imperfect, suggests clustering of phenotype-genotype correlation.

To date there is no cure for *STAT1*-related disorders. Although control of bacterial infections can be achieved

Available online 11 September 2014

LymphoSign Journal 1:57–59 (2014)
dx.doi.org/10.14785/lpsn-2014-0011



Figure 1: Typical nail changes found in patients with CMCC due to mutations in *AIRE* or in *STAT1*, as indicated.




Figure 2: Mean (black line) and median (red line) of the age of 24 reported cases with mutations in the coiled-coil domain (CCD) and the 13 cases with mutations in the DNA binding domain (DBD). Patients with mutations in the DBD were reported at a significantly ($p = 0.00004$) younger age than patients with mutations in the CCD.

by using antibiotic therapy and immunoglobulin replacement, other manifestations cannot be prevented or effectively treated. Attempts to clear chronic cytomegalovirus, Epstein–Barr virus, or John Cunningham virus




Figure 3: Mean (black line) and median (red line) of reported age of deaths of 10 cases with mutations in the coiled-coil domain (CCD) and 6 cases with mutations in the DNA binding domain (DBD). Patients with mutations in the DBD die at a significantly younger age ($p < 0.005$) than patients with mutations in the CCD.

were not successful in these patients. Moreover, vascular changes to the central nervous system, including vasculitis, aneurisms, and acute bleeding cannot be effectively and completely prevented, even with harsh immunosuppressive treatments with cyclophosphamide. The use of high dose intravenous immunoglobulin failed to show a significant improvement, but the use of immune modulation agents such as Mycophenolate Mofetil may be of benefit in controlling some autoimmune features.

For cases with mutations in the DBD domain, especially those known to be associated with mortality at an early age, such as T385M, hematopoietic stem cell therapy might be a plausible consideration.

REFERENCES

Ahonen, P., Myllärniemi, S., Sipilä, I., and Perheentupa, J. 1990. Clinical variation of autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED) in a series of 68 patients. *N. Engl. J. Med.* **322**:1829. PMID: 2348835. doi: 10.1056/NEJM 199006283222601.

Bentur, L., Nisbet-Brown, E., Levison, H., and Roifman, C.M. 1991. Lung disease associated with IgG subclass deficiency in chronic mucocutaneous candidiasis. *J. Pediatr.* **118**:82. PMID: 1986107. doi: 10.1016/S0022-3476(05)81852-9.

Finnish–German APECED Cosortium. 1997. An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PHD-type zinc-finger

- domains. *Nat. Genet.* **17**:399. PMID: 9398840. doi: 10.1038/ng1297-399.
- Liu, L., Okada, S., Kong, X.F., Smeekens, S.P., Joosten, L.A.B., Gilissen, C., Arts, P., Rosentul, D.C., Carmichael, A.J., Smits-van der Graaf, C.A.A., Bart Jan Kullberg, B.J., van der Meer, J.W.M., Desa Lilic, D., Veltman, J.A., and Netea, M.G. 2011. Gain-of-function human *STAT1* mutations impair IL-17 immunity and underlie chronic mucocutaneous candidiasis. *J. Exp. Med.* **208**:1635. PMID: 21727188. doi: 10.1073/pnas.0501063102.
- Maródi, L., Cypowwyj, S., Tóth, B., Smeekens, S.P., Joosten, L.A.B., Gilissen, C., Arts, P., Rosentul, D.C., Carmichael, A.J., Smits-van der Graaf, C.A.A., Bart Jan Kullberg, B.J., van der Meer, J.W.M., Desa Lilic, D., Veltman, J.A., and Netea, M.G. 2012. Molecular mechanisms of mucocutaneous immunity against *Candida* and *Staphylococcus* species. *J. Allergy. Clin. Immunol.* **130**:1019. PMID: 23040277. doi: 10.1016/j.jaci.2012.09.011.
- Sampaio, E.P., Hsu, A.P., Pechacek, J., Bax, H.I., Dias, D.L., Paulson, M.L., Chandrasekaran, P., Rosen, L.B., Carvalho, D.S., Ding, L., Vinh, D.C., Browne, S.K., Datta, S., Milner, J.D., Kuhns, D.B., Long Priel, D.A., Sadat, M.A., Shiloh, M., De Marco, B., Alvares, M., Gillman, J.W., Ramarathnam, V., de la Morena, M., Bezrodnik, L., Moreira, I., Uzel, G., Johnson, D., Spalding, C., Zerbe, C.S., Wiley, H., Greenberg, D.E., Hoover, S.E., Rosenzweig, S.D., Galgiani, J.N., and Holland, S.M. 2013. Signal transducer and activator of transcription 1 (*STAT1*) gain-of-function mutation and disseminated coccidioidomycosis and histoplasmosis. *J. Allergy. Clin. Immunol.* **131**:1624. PMID: 23541320. doi: 10.1016/j.jaci.2013.01.052.
- Sharfe, N., Nahum, A., Newell, A., Dadi, H., Ngan, B., Pereira, S.L., Herbrick, J.A., and Roifman, C.M. 2014. Fatal combined immunodeficiency associated with heterozygous mutation in *STAT1*. *J. Allergy. Clin. Immunol.* **133**:807. doi: 10.1016/j.jaci.2013.09.032.
- Takezaki, S., Yamada, M., Kato, M., Park, M.J., Maruyama, K., Yamazaki, Y., Chida, N., Ohara, O., Kobayashi, I., and Ariga, T. 2012. Chronic Mucocutaneous candidiasis caused by a gain-of-function mutation in the *STAT1* DNA-binding domain. *J. Immunol.* **189**:1521. PMID: 22730530. doi: 10.4049/jimmunol.1200926.
- Uzel, G., Sampaio, E.P., Lawrence, M.G., Hsu, A.P., Hackett, M., Dorsey, M.J., Noel, R.J., Verbsky, J.W., Freeman, A.F., Janssen, E., Bonilla, F.A., Pechacek, J., Chandrasekaran, P., Browne, S.K., Agharahami, A., Gharib, A.M., Mannurita, S.C., Yim, J.J., Gambineri, E., Torgerson, T., Tran, D.Q., Milner, J.D., and Holland, S.M. 2013. Dominant gain-of-function *STAT1* mutations in FOXP3 wild-type immune dysregulation-polyendocrinopathy-enteropathy-X-linked-like syndrome. *J. Allergy. Clin. Immunol.* **131**:1611. PMID: 23534974. doi: 10.1016/j.jaci.2012.11.054.
- Van de Veerdonk, F.L., Plantinga, T.S., Hoischen, A., Smeekens, S.P., Joosten, L.A.B., Gilissen, C., Arts, P., Rosentul, D.C., Carmichael, A.J., Smits-van der Graaf, C.A.A., Bart Jan Kullberg, B.J., van der Meer, J.W.M., Desa Lilic, D., Veltman, J.A., and Netea, M.G. 2011. *STAT1* mutations in autosomal dominant chronic mucocutaneous candidiasis. *N. Eng. J. Med.* **365**:54. doi: 10.1056/NEJMoa1100102.

Chaim M. Roifman, MD, FRCP, FCACB
Editor-in-Chief
(croifman@lymphosign.com)